

SLURRY FOR SOIL™ HC

QUICK START GUIDE

INSTRUCTIONS:

Preparing SlurryForSoilHC is very quick and easy:

STEP 1: Heat water to approximately 37°C. Add about 5L of warm water per kg of SlurryForSoil.

STEP 2: Stir well until the powder dissolves and leave for at least 45min.

STEP 3: Stir solution again, before adding to the liquid layer of the slurry.

IMPORTANT USER NOTES:

- Step 1 - water temperature – activating water temperatures can range from 30°C to 37°C. Do not exceed 40°C as you will destroy some of the microbes.
- Step 2 – while SlurryForSoil contains a limited food source, soon after activation the microbes will need more. The solution should be added to slurry ideally 45 mins to an hour after mixing but no longer than 16 hours after mixing.
- Step 3 – it's important to add the solution to the liquid fraction of slurry and not the sludge at the bottom, nor through the crust at the top. If possible, deliver directly, or through the slats or chutes. For large underground tanks, apply in 2 or 3 places.
- Do not stir the slurry once SlurryForSoil has been added. The bacteria and fungi need to be concentrated in order to spread and out-grow and dominate existing microbes present in the slurry
- Avoid contact with eyes and avoid inhaling product when mixing. Wash hands after use. Always read the label and product information before use.
- Once treated, leave slurry for at least 6-8 weeks and apply a booster dose after 12 weeks.

FURTHER INFORMATION:

Contact us at info@sylgenanimalhealth.com
or call +44 (0)1559 370222

Get the latest news at www.sylgenanimalhealth.com
and on social media:

